

## Differences between ethnic minority and native children in breakfast habits

*Giorgia Monti<sup>1</sup>, Dimitra Gkliati<sup>1</sup>, Lavinia Fanciullo<sup>1</sup>, Carla Mastroianni<sup>1</sup>, Viviana Finistrella<sup>2</sup>, Elio Volta<sup>2</sup>, Anna Ruggerini<sup>2</sup>, Roberto Ciati<sup>2</sup>, Maurizio Vanelli<sup>1,2</sup>*

<sup>1</sup>Postgraduate School of Paediatrics University of Parma, and <sup>2</sup>Giocampus Scientific Committee, Parma, Italy

**Abstract.** Aim of this study was to investigate the breakfast habits in a cohort of school ethnic minority (EM) children admitted to a summer sport school (SSS) in order to evaluate the possible differences with their Italian peers (IP). One hundred-seventy-nine children aged 6-14 years were interviewed through a questionnaire about breakfast behaviors and nutrition habits. The results were compared with those obtained in a randomized sample of 179 Italian peers (IP) attending the same SSS. EM children skip breakfast more frequently than IP (15% vs 7%;  $p=0.029$ ). The omissions are attributed to the lack of time (45% vs 4%) and not being hungry upon waking (44% vs 38%). During weekends the percentage of breakfast skippers decreased of about 3 percentage points but persisted a significant difference between the two groups ( $p=0.024$ ). EM children who did not have breakfast show a mean BMI higher than breakfast consumers ( $p=0.05$ ). Bakery products and milk were the most frequently consumed food and drink in EM and IP children (68% vs 70% and 71% vs 76% respectively). EM children chose and prepared personally breakfast at home more frequently than IP (54% vs 22%). These EM children were more inclined to skip breakfast ( $p=0.025$ ) and to have an elevated BMI ( $p=0.006$ ) than IP. An educational program, for ethnic minority families, in different languages and according to their cultural, religious and socio-economic influences could be a possible means for changing their lifestyle and reducing their risk to become overweight. ([www.actabiomedica.it](http://www.actabiomedica.it))

**Key words:** ethnic minority, breakfast, obesity, overweight, food, nutrition

### Introduction

Immigration is a growing phenomenon in the developed countries (1). In Italy, the amount of migrant people is estimated in about 7% of the whole population (2). Many habits of these ethnic minorities (EM) are producing demographic and health problems in the Occidental World. Immigrant birth rate is higher than that of natives (3). Immigrated children face many different health risks when compared to native children: they are subjected to many infectious diseases no longer seen in our countries such as malaria, tuberculosis, and HIV; they are more likely to have inadequate immunity to vaccine-preventable illnesses;

and finally they show a higher likelihood of having malnutrition and developmental delay (4).

Diets of immigrated children worsen the longer they stay in developed countries (4), but their real food habits have been rarely investigated (5, 6). Some studies highlighted that children and adolescents belonging to an EM have a high risk to become overweight or obese. This risk seems to be related to a poor food quality consumption, a low socioeconomic and cultural parental level (4, 7, 8), and to a high susceptibility to environmental factors (9).

Family environment seems to have an important role on children and adolescent dietary mistakes (10-12). A most frequent mistake in childhood concerns

the skipping breakfast. We previously found that this habit occurred in 22% of a native youth population, and it was strictly related to overweight and obesity (13). It is well known that people who skip breakfast are prone to snack more frequently junk foods than natural snacks such as fruits and vegetables (14-16).

Reports on breakfast habits in ethnic minority children are generally unknown. In this paper we have investigated breakfast habits in a cohort of EM children participating to a summer sport school in order to evaluate the possible differences with their Italian peers (IP).

## Materials and methods

### *Subjects*

Data for the present study came from the participants in "Giocampus", a Summer Sport School for children, aged 6 to 14 years, which took place in the Campus sport area of the University of Parma from June to September 2010, in collaboration with Postgraduate Medical School of Paediatrics, Graduate School of Sport and Exercise Sciences of the University of Parma, Town Council, School Inspectorate, University Sport Club, and Barilla Group. The children admitted to this Summer Sport School, spent 8 hours per day from Monday to Friday in physical activities (i.e. soccer, volleyball, basketball, artistic dance, tennis, fencing and athletics) under the supervision of professional instructors. During this time, snacks, lunch and beverage were regularly distributed. Breakfast was not included.

Informed consent and assent for participation in the study were obtained from parents and children respectively. For privacy reasons, education level was not recorded. The survey was approved by the Ethical Committee of the University of Parma.

### *Procedure of interview and growth evaluation*

All the participants were requested to answer a questionnaire on their breakfast habits. The questionnaire has been described in a previous report (13). Children were randomly recruited and finally divided

into 2 homogeneous groups about their ethnic origin. Children were asked whether, when, where, how and with whom they consumed breakfast during weekdays and at the weekend; who prepared meals; what they ate and drank; what they did during breakfast. Breakfast was defined as any intake of food or beverage between 6 and 8 a.m. before going to the Summer Sport School. The reports not eating a morning meal at home, or breakfast, fewer than three times per week were defined: "Skip breakfast".

Harpندن stadiometers were used for height, and every child was measured according to the standard technique elsewhere described (17). Body mass index (BMI) was calculated using the formula: weight (kg)/height (m<sup>2</sup>). Overweight and obesity were defined according to the international cut off points for body mass index for overweight and obesity proposed by Cole et al. (18).

### *Statistical analyses*

The data collected were analyzed using SPSS for Windows and expressed as mean  $\pm$  SD. Numerical differences were calculated by Student's t-test. The  $\chi^2$  test was used to study the differences between proportions. Differences were considered significant if  $p < 0.05$ .

## Results

The children who accepted to answer the questionnaire were globally 364 (37.6% of all participants in Giocampus), 228 were boys and 336 girls: 179 children came from an EM (Group 1) and 185 children were native (Group 2). All families of EM children originated from Sub Saharan Countries and have been living in Parma since long. The base-line characteristics of the study cohorts are summarized in Table 1.

EM children were prone to skip breakfast more frequently than the IP children (15% vs 7%;  $\chi^2=4.756$   $p=0,029$ ). During weekends the percentage of breakfast skippers decreased in both groups to 12% and to 5% respectively, but the difference between groups (12% vs 5%) persisted significant ( $\chi^2=5.086$   $p=0.024$ ). The omissions of breakfast were attributed to the lack

**Table 1.** Baseline characteristics of the participants in the study

	Group 1 (Ethnic group)	Group 2 (Native group)	p
No. of children	179	185	
Age (yr)	9.47±2.02	9.46±2.01	0.963
Male sex (%)	64%	64%	
BMI	18.11±3.12	18.25±2.99	0.970

time (45% vs 54%;  $p=0.258$ ) and not being hungry upon waking (44% vs 38%;  $p=0.472$ ) in 1<sup>st</sup> and 2<sup>nd</sup> Group respectively.

Breakfast skipper children in Group 1 had a mean BMI higher (19.19±3.5) than those who regularly ate breakfast in Group 2 (17.92±3.01;  $t=1.970$   $p=0.050$ ).

Thirty four percent of EM and 27% percent of IP children reported to have breakfast alone ( $p$  0.155). These EM children resulted to have a mean BMI (18.79±2.98) higher than children who consumed breakfast with their parents (17.64±3.03;  $p=0.018$ ) without differences between groups.

Fifty-four percent of EM children reported to prepare breakfast themselves more frequently than IP children (22%;  $p=0,000$ ). Children who personally chose and prepared breakfast were inclined to skip breakfast both in group 1 (20% vs 8%;  $\chi^2=5.025$ ;  $p=0,025$ ) and in group 2 (13% vs 4%;  $\chi^2=4.114$ ;  $p=0,043$ ).

Twenty-four percent of children in Group 1 and 28% of children in Group 2 referred to watch TV during breakfast .

Bakery products were the most frequently consumed as breakfast foods the Groups 1 and 2 (68% and 70% respectively). Fruit was in general sporadically eaten, but EM girls 6 to 10 years old ate fruit more than IP (16% vs 6%;  $p=0.042$ ). Milk with or without chocolate was the most popular breakfast drink in both EM and IP children groups (71% and 76%) (Table 2).

## Discussion

It is proved that starting each day with breakfast improves the school performance, guarantees an adequate intake of energetic nutrients, and reduces the risk of overweight and obesity (16, 19-21). Several

**Table 2.** Frequency of daily consumption of different foods and drinks at breakfast in the two ethnic groups of children

Foods and Drinks	Boys %		Girls %	
	Native	Ethnic	Native	Ethnic
Biscuits	32	34	34	32
Snack cakes	16	17	17	20
Cereals	13	10	8	16
Crackers	6	9	4	7
Cakes	5	5	4	8
Bread	7	4	11	6
Butter and marmalade	4	5	5	4,5
Fruit	10	9	13	6
Milk	54	52	48	53
Milk and cocoe	22	23	14	25
Tea	6	11	21	4
			T=11.703 p=0.000	
Fruit Juices	12	11	11	17
Hot chocolate	5	2	1	

studies have consistently reported positive association between adiposity in children and skipping breakfast (14, 22-24).

The percentage of children which do not have breakfast usually ranges from 10% to 30% depending on the age, gender, ethnic group and country of origin (20). In Italy a recent study evidenced that about 11% of children skips breakfast (25). In Parma, in 2005, we found an unexpected high number of skippers in native children (22%) (13).

Detailed observations on breakfast habits in immigrant children are not frequently found in literature (6). Anecdotic reports highlight an upward trend in young immigrant people to avoid breakfast and to become obese in comparison with native children (6). In the present study we have found that EM children are prone to skip breakfast (15%) more frequently than their native peers (7%). This bad habit exposed EM children to a higher risk ( $p=0.05$ ) to become overweight or obese. The reasons of this omission are consistent with those previously described: lack of time, and not being hungry upon waking (12, 26, 27). It well known that 70% of primary school children in Italy are inclined to go to bed late at around 10:00 p.m. (13). This habit might lead children to wake up

late and to skip breakfast in order to be on time for school.

It is encouraging to observe also in the present study that during weekends the percentage of skippers breakfast decline both in EM and IP children. This improvement has been attributed to the presence of parents at home which thus can personally manage and take breakfast together with their offspring (13). Anyway, the skippers in EM group continue to be more numerous than in native group also in weekends. We can speculate that in EM group this happens because many parents work during weekend too.

Parental model is considered as an important factor to stimulate children to have breakfast. Low socio-economic and cultural level, and the discontinuous presence of the parents at home affect this target especially in the ME group (28, 29). It is known that children which are left alone at home and are responsible for preparing their own meal are more frequently inclined to skip breakfast (13). This phenomenon is difficult to be solved in ME children because important social and economic reasons come into play.

Bakery products (biscuits, snacks, cereals) and milk were the favorite foods and drinks in both children groups. Under this point of view EM children showed to be completely integrated in the social behavior. It is nevertheless remarkable to observe that Italian young girls prefer to drink fruit juices than milk, while their ethnic minority peers choose rather milk and tea. Increasing in preference for the fruit juices has been recently observed also in American teenagers without distinction between ethnic origin (30).

In conclusion, immigrant children skip breakfast more frequently than IP and have a higher risk to become overweight and obese. The absence of parents at home and the inappropriate role of EM children in the choice of breakfast foods chiefly contribute to this incorrect diet behavior. Face to these habits and related metabolic consequences, it is urgent to find some corrective interventions. The first one could concern the possibility to recover at school the breakfast neglected at home. A similar option has been evaluated in previous "Giocampus" editions with a great satisfaction of both EM children and their parents. The second intervention is to carry out a convincing educational program in the EM families using booklets

and TV spots in different languages. In this perspective, a Multilanguage booklet, titled "Alarm obesity", has been already placed available in "Giocampus" web site. Breakfast promotion in Mosques is not to be finally excluded.

## Acknowledgements

Thanks go to doctor Rossana Di Marzio for her support in the preparation of this manuscript

## References

1. Ministero dell'Interno. 1° rapporto sugli immigrati in Italia. Dicembre 2007. [www.interno.it](http://www.interno.it)
2. ISTAT. Popolazione straniera residente in Italia. [www.istat.it](http://www.istat.it)
3. Camarota S A. Birth rates among immigrants in America comparing fertility in the U.S. and home Countries. [www.cis.org/articles/2005](http://www.cis.org/articles/2005)
4. <http://www.journals.elsevierhealth.com/periodicals/yspid/article/PIIS1045187004001256/ab>
5. Cornelisse-Vermaat JR, Maassen van den Brink H. Ethnic differences in lifestyle and overweight in The Netherlands. *Obesity* 2007; 15: 483-93.
6. Brussaard JH, Van Erp-Baart MA, Brants HAM, et al. Nutrition and health among migrants in The Netherlands. *Public Health Nutr* 2001; 4: 659-64.
7. Cullen KW, Baranowski T, Owens E, et al. Ethnic differences in social correlates of diet. *Health education research* 2001; 17: 7-18.
8. Hulshof KFAM, Brussaard JH, Kruijzinga AG, et al. Socio-economic status, dietary intake and 10 y trends: the Dutch National Food Consumption Survey. *Eur J Clinical Nutr.* 2003; 57: 128-37.
9. Swinburn BA, Egger GJ, Raza F. Dissecting obesogenic environments: the development and application of a framework for identifying and prioritizing environmental interventions for obesity. *Preventative Med* 1999; 29: 563-70.
10. Patrick H, Nicklas TA. A review of family and social determinants of children's eating patterns and diet quality. *J Am College Nutrition* 2005; 24 (2): 83-92.
11. Pearson N, Biddle SJH, Gorely T. Family correlates of breakfast consumption among children and adolescents. A systematic review. *Appetite* 2009; 52: 1-7.
12. Harding S, Teyhan A, Maynard MJ, et al. Ethnic differences in overweight and obesity in early adolescence in the MRC DASH study: the role of adolescent and parental lifestyle. *Int J Epidemiol* 2008; 37: 162-72.
13. Vanelli M, Iovane B, Bernardini A, et al. Breakfast habits of 1,202 northern Italian children admitted to a summer sport school. Breakfast skipping is associated with overweight and obesity. *Acta Biomed* 2005; 76: 79-85.

14. Meropi D, Kontogianni MD, Farmaki AE, et al. Associations between lifestyle patterns and Body Mass Index in a sample of Greek children and adolescents. *J Am Diet Assoc* 2010; 110: 215-21.
15. Vereecken C, Dupuy M, Rasmussen M, Kelly C, et al. Breakfast consumption and its socio-demographic and lifestyle correlates in schoolchildren in 41 countries participating in the HBSC study. *Int J Public Health* 2009; 54: S180-S190.
16. Breakfast in America: The National Health and Nutrition Examination Survey (NHANES), 2001-2002. [http://www.ars.usda.gov/SP2UserFiles/Place/12355000/pdf/Breakfast\\_2001\\_2002.pdf](http://www.ars.usda.gov/SP2UserFiles/Place/12355000/pdf/Breakfast_2001_2002.pdf).
17. Cameron M. The methods of auxological anthropometry. In Falkner F, Tanner JM eds: *Human Growth. A comprehensive treatise*, 2nd ed. Plenum press, New York, 1986: 3-46.
18. Cole TJ, Bellizzi C, Flegal KM, et al. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* 2000; 320: 1240-3.
19. Marangoni F, Poli A, Agostini C, et al. A consensus document on the role of breakfast in the attainment and maintenance of health and wellness. *Acta Biomed* 2009; 80: 166-71.
20. Rampersaud GC, Pereira MA, Girard BL, et al. Breakfast habits, nutritional status, body weight, and academic performance in children and adolescents. *J Am Diet Assoc* 2005; 105: 743-60.
21. Dubois L, Girard M, Potvin Kent M, et al. Breakfast skipping is associated with differences in meal patterns, macronutrient intakes and overweight among pre-school children. *Public Health Nutr* 2008; 18: 1-10.
22. Hansen K, Joshi H. Millennium Cohort Study Third Survey: a user's guide to initial findings, p. 162. London: Centre for Longitudinal Studies, Institute of Education 2008.
23. Haug E, Rasmussen M, Samdal O, et al. Overweight in school-aged children and its relationship with demographic and lifestyle factors: results from the WHO-Collaborative Health Behaviour in School-aged Children (HBSC) Study. *Int J Public Health* 2009; 54 (2): 167-79.
24. Croezen S, et al. Skipping breakfast, alcohol consumption and physical inactivity as risk for overweight and obesity in adolescents: results of the E-MOVO project. *Eur J Clin Nutr* 2009; 63: 405-12.
25. Spinelli A, Lamberti A, Baglio G, et al. OKkio alla SALUTE: sistema di sorveglianza su alimentazione e attività fisica nei bambini della scuola primaria. Risultati 2008. Roma: Istituto Superiore di Sanità; 2009 (Rapporti ISTISAN 09/24).
26. Sweeney N, Horishita N. The breakfast-eating habits of inner city high school students. *J Sch Nurs* 2005; 21: 100-5.
27. Reddan J, Wahlstrom K, Reicks M. Children's perceived benefits and barriers in relation to eating breakfast in schools with or without Universal School Breakfast. *J Nutr Educ Behav* 2002; 34: 47-52.
28. Hanson MD, Chen E. Socioeconomic status and health behaviors in adolescence: a review of the literature. *J Behavioral Med* 2007; 30: 263-85.
29. Delva J, O'Malley PA, Johnston LD. Racial/ethnic and socioeconomic status differences in overweight and health-related behaviors among American students: national trends 1986-2003. *J Adolesc Health* 2006; 39: 536-45.
30. Evans A.E, Springer A.E, Evans M.H, et al. A descriptive study of beverage consumption among an ethnically diverse sample of public school students in Texas. *J Am College Nutrition* 2010; 29: 387-396.

---

Accepted: May 21th 2011

Correspondence: Giorgia Monti, MD  
 Postgraduate School of Paediatrics  
 University of Parma, Italy  
 v.le A. Gramsci, 14  
 43100 Parma, Italy  
 E-mail: giorgia-monti@alice.it