

Screening for antioxidant capacity, pollen types and phytochemical profile by GC/MS and UHPLC from propolis

Nesrin Ecem Bayram^{1,2*}, Yusuf Can Gerçek³, Gül Cevahir Öz³

¹ Department of Food Processing, Aydıntepe Vocational School, Bayburt University, Bayburt, Turkey - E-mail address: ecem.nesrin@gmail.com; ² Bayburt University Beekeeping Research, Development and Application Center, Bayburt, Turkey; ³ Department of Biology, Faculty of Science, Istanbul University, Istanbul, Türkiye

Summary. Propolis is a resinous mixture collected by honeybees from different parts of plants such as poplars, birches, alders, conifers, pines, palms and willows. This study aims to determine pollen types (by light microscope), chemical compound profile (by GC-MS), fatty acid composition (by GC-MS), total phenolic content (by the Folin-Ciocalteu method), total flavonoid content (by the aluminium chloride method) and antioxidant capacity (by the CUPRAC, ABTS and CERAC methods) of a propolis sample from the western part of Istanbul, Turkey. As a result of microscopic analysis of the sample, pollen types of taxa belonging to 27 plant families were diagnosed. The GC-MS analysis of propolis revealed the presence of 38 phytochemical constituents that may contribute to its quality. Of these compounds, rates of “4H-1-benzopyran-4-one, 2,3-dihydro-5,7-dihydroxy-2-phenyl-,(S)” were highest. In addition, the concentrations of naringenin, pinocembrin and galangin were determined to be 2.45, 9.92 and 7.06 mg/ml by UHPLC analysis. The extract had significant antioxidant activity in all assays, with values of 282.8±9 mg TE /g in the CUPRAC, 425.7±18 mg TE/g in the CERAC and 186.4±8 mg TE/g in the ABTS assays. Antioxidant capacity of the propolis extract was positively associated with the total phenolic and flavonoid contents of the extract. Moreover, the major fatty acids were C20:1n9 (cis-11-eicosenoic acid), C22:1 (erucic acid) and C24:1 (nervonic acid). The results show that the propolis used in our study has important potential as an alternative food supplement and for cosmetic and therapeutic medicine and it can be used as an active agent in these areas.

Keywords: Propolis, pollen content of propolis, fatty acid of propolis, antioxidant activity.

Introduction

Propolis (bee glue) is a natural resin-like substance collected by honeybees. The chemical composition of propolis is very complex. To date, more than 500 chemical components have been identified in its chemical structure (1, 2). The main components of propolis are pollen (5%-10%), oil and wax (30%-50%), resin (50%-70%), and other chemical compounds including: vitamins B, C and E, sugars, minerals, phenols, flavonoids, amino acids, fatty acids, steroids and stilbenes as well as aromatic compounds (3). The diver-

sity and rate of these constituent's change depending on the region of collection, plant source and weather conditions when it was collected.

Propolis has been used by people for therapeutic purposes for centuries. It has antioxidant (4), antibacterial (5) and antifungal (6), antiviral (7), antiinflammatory (8), antitumor, hepatoprotective, antipsoriatic (9), antihyperalgesic (10) and antigenotoxic (11) effects. In particular, several studies have reported that flavonoids from the main components of propolis are responsible for the majority of these activities (12-14) The degree of these effects varies according to the ac-

tive substances contained in the propolis and the contents of the propolis samples produced in different regions also varies, which makes standardization difficult. Therefore, we evaluated the phenolic and chemical profile in addition to the antioxidant capacity of propolis from İstanbul (Turkey) in order to contribute to standardization efforts in this study. We also examined the fatty acid composition.

Material and Methods

Chemicals

Trolox (6-hydroxy-2,5,7,8-tetramethylchroman carboxylic acid), 2,20 -Azino-bis(3-ethylbenzothiazoline-6-sulphonic acid), gallic acid, quercetin, sodium carbonate, Folin-Ciocalteu's phenol reagent and gallic acid were purchased from Sigma-Aldrich (Istanbul). Cu (II) chloride, neocuproin, ammonium acetate, cerium sulphate, aluminium chloride, sodium hydroxide, sodium nitrite and ethanol were obtained from Merck. All chemicals used in the experiments were of analytical grade.

Geographical origin of propolis

A propolis sample was collected from an apiary located in the western part of İstanbul, Turkey in 2018. It was then pulverized with a grinder and stored in the freezer (-18°C) until analysis.

Microscopic investigation of propolis

The pollen spectra of the propolis sample was detected according to the methodology of Warakomska and Maciejewicz (1992) (15). 1 g of the propolis sample was mixed with ethanol-ether-acetone (1:1:1) and vortexed. This mixture was filtered through a strainer with 0.3 mm holes. The suspension was centrifuged for 20 minutes at 3500-4000 rpm. The supernatant liquid was then poured off; glycerin gelatin with basic fuchsin about 1-2 mm³ in width was taken with a sterile needle and transferred onto a slide after imbruing it in the pellet in the bottom of the centrifuge tube. The slide in this form was heated at 30-40°C to allow the dissolution of fuchsin glycerin gelatin; and then covered with a 18x18 lamella. Then, it was examined with a Leica DM500 light microscope (16).

Preparation of propolis extract

The process of extraction from propolis by ultrasound treatment was performed using a 25 kHz ultrasonic processor (model VCX 750; Sonics & Materials, Inc., Newtown, CT, USA). Thirty grams of propolis powder was dispersed with 100 mL of ethanol (LC-grade). Sonication was carried out in a double-walled stainless steel chamber using a titanium probe with an emitting face 19 mm in diameter that was kept immersed 2 cm below the surface of the sample during sonication. The temperature was kept constant throughout sonication by circulating water through the jacket of the chamber. The samples were exposed to extract at 40% amplitude for 5 minutes at 30°C. The extract was cooled to room temperature and filtered through Whatman no 1 filter paper and 0.22 µm polypropylene filter, then transferred to amber bottles and stored at -18°C until analysis.

Chemical screening of propolis composition by GC-MS

The filtered solution was diluted in 1:10 ratio (w/v) with 96% ethanol and evaporated to complete dryness. About 5 mg of the dry matter was mixed with 75 µL of dry pyridine and 50 µL bis (trimethylsilyl) trifluoroacetamide (BSTFA) and heated at 80°C for 20 min. The final supernatant was analyzed by gas chromatography-mass spectroscopy (GC-MS). The extract was analyzed using a GC 7890A from Agilent (Palo Alto, CA, USA) coupled with mass detector (MS 5975C, Agilent) equipped with a DB-5 MS capillary column (30 m × 0.25 mm and a film thickness of 0.25 µm). The column oven temperature was initially maintained at 50°C for 1 min, then programmed to rise to 150°C at 10°C/min and maintained for 2 min. Finally, the temperature was increased to 280°C at 20°C/min and maintained at 280°C for 5 min. Helium was used as the carrier gas at a flow rate of 1 mL/min. For GC-MS detection, an electron ionization system with ionization energy of 70 eV was used. The mass spectra were recorded within 40-550 (m/z), full scan mode, that revealed the total ion current (TIC) chromatograms. MS transfer line temperatures were set at 265°C (2).

The active ingredients (naringenin, pinocembrin and galangin) of propolis by UHPLC

In the study, double distilled water and formic

acid (19: 1, v: v) were used as the mobile phase and the separation of phytochemicals was performed on a reverse phase column Perkin Elmer RP-18 (USA) (12.5 x 0.4 cm, 5 µm particle size). Formic acid and water (A) and methanol (B) have a constant solvent flow rate of 1 mL of min⁻¹, starting isocratically with 30% B in A for up to 15 minutes, then loading a gradient to reach 40% B in 20 minutes, 45% B at 30 minutes, 60% B at 50 minutes, 80% B at 52 minutes and 90% B at 60 minutes, then isocratic for 65 minutes. The propolis extract was analyzed using a UHPLC Flexar using a multi-channel photodetector FX-20 and samples were injected with a Flexar autosampler FX-15 device. The column was maintained at room temperature and the chromatograms were eluted with Chromera Manager software (Perkin Elmer, USA).

Bioactivity tests

Total phenolic content (TPC)

The content analysis of total phenolic compounds was performed to the Folin-Ciocalteu method proposed by Magalhães et al. (2010) using gallic acid as a reference standard (17). 50 µL of sample and 50 µL of Folin-Ciocalteu reagent (1:5, v/v) were placed in each well. After that, 100 µL of sodium hydroxide solution (0.35 M) was added. The absorbance at 760 nm of the blue complex formed was read after 3 minutes. The results were expressed as gallic acid equivalent (mg GAE/g).

Total flavonoid content (TFC)

Total flavonoid analysis of the extract was performed using a modified version of the method Jia et al. (1999) used in their work (18). Accordingly, an aliquot (1 mL) extract was mixed with 0.3 mL 10% AlCl₃.6H₂O solution after the addition of 0.3 mL 5% NaNO₂ solution. 2 mL of 1 M NaOH solution was added and 2.4 mL of water was added and the mixture was stirred. At 510 nm the absorbance was measured against the prepared reagent blank by Epoch Multi-Detection Microplate Reader with 96-well plates (BioTek Instruments, Inc., P). Total flavonoid content was expressed as mg quercetin equivalent (mg QE/g).

Determination of antioxidant activity

Cupric ion reducing antioxidant capacity (CUPRAC) assay: To a test tube was added 1 mL of 1.0 x

10⁻² M Cu (II) chloride solution, 1 mL of 7.5 x 10⁻³ M neocuproin solution and 1 mL of 1 M ammonium acetate buffer (pH 7.0), respectively. Then x mL propolis extract was stirred into 95% ethanol-water (1.1-x). The tube was kept at room temperature for 30 minutes. At the end of the period, the absorbance value of the solution at 450 nm was measured against the non-including antioxidant solution by Epoch Multi-Detection Microplate Reader with 96-well plates (BioTek Instruments, Inc., P) (19).

Modified cerium (IV)-based antioxidant capacity (CERAC) assay: The total antioxidant capacity of the tested sample was also determined using the Cerium (IV) assay of Ozyurt et al. (2007). A 1.0 mL 2.0 x 10⁻³ M Ce(SO₄)₂ + x mL sample + (9-x) mL H₂O solution with a total volume of 10.0 mL is prepared and left at room temperature for 30 minutes. After 320 nm the absorbance is measured against the distilled water and compared to a Trolox standard curve of 1-20 mM (20).

2,2'-azino-bis-3-ethylbenzthiazoline-6-sulphonic acid (ABTS) assay: The ABTS reagent prepared at 7 mM concentration was dissolved in water. The ABTS radical cation was reacted with 2.45 mM potassium persulfate and left in the dark at room temperature for 12-16 hours before use. To study the sample, the ABTS solution was diluted with ethanol to an absorbance of 0.70 at 734 nm and equilibrated at 30°C. 1 ml of sample and 1 ml of ABTS solution, were diluted with methanol to a total volume of 4 ml. After the tube was kept closed at room temperature for 6 minutes, the absorbance value of the sample was read at 734 nm (21).

Fatty acid methyl ester (FAME) analysis of propolis by GC-MS

Lipid extraction

2 g of pulverized raw propolis was weighed into a soxhlet cartridge, which was placed in the extraction chamber in the soxhlet apparatus (Buchi B-811). A 100 mL aliquot of n-hexane was transferred into the solvent cup and placed on the heating plates. The cooling water supply to the condensers was opened to ensure continuous recycling of the solvent and temperature selected as per the Büchi manual for extraction in the continuous mode.

Preparation of FAME

A 100 mg sample (propolis oil) was weighed in a 20 mL test tube and dissolved in 10 mL of n-hexane. 0.1 mL of 2N KOH solution was added to tube. This mixture was vortexed for 30 seconds and then centrifuged. 2 mL of saturated NaCl solution was added and the organic phase was separated. 1 µL of the final solution was analyzed in GC-MS (22).

FAME Analysis by GC-MS

Methylated fatty acid samples were analyzed by Agilent 6890 GC gas chromatography and 5973 MSD mass spectrometry. DB-23 60 m x 0.25 mm ID, 0.15 µm (J&W 122-2361) column was used in the analyses. Helium was used as carrier gas. The oven temperature was initially maintained at 50°C for 1 min and then programmed to rise to 175°C at 25°C/min. Finally, the temperature was increased to 230°C at 4°C/min and maintained at 280°C for 5 min. The injection temperature was set to 230°C. 1 µL injection was made and the split ratio was adjusted to 1/50 (23).

Results and Discussion

The chemical structure of propolis is quite complex and varies considerably depending on environmental factors, flora and vegetation of the region. While bees collect the propolis from different plant sources, they also enrich its chemical structure by adding plant pollen, and the chemical composition of propolis contains around 5% pollen. Pollen types of the Aceraceae (<1%), Apiaceae (7.75%), Asteraceae (16.18%), Boraginaceae (4.55%), Brassicaceae (2.69%), Campanulaceae (2.69%), Caprifoliaceae (<1%), Caryophyllaceae (2.86%), Cyperaceae (<1%), Dipsacaceae (<1%), Fabaceae (19.56%), Fagaceae (7.08%), Geraniaceae (<1%), Lamiaceae (16.52%), Malvaceae (<1%), Onagraceae (<1%), Papaveraceae (1.34%), Plantaginaceae (<1%), Poaceae (4.34%), Polygonaceae (1.01%), Pinaceae (<1%), Ranunculaceae (<1%), Rosaceae (1.34%), Liliaceae (<1%), Rubiaceae (1.01%), Salicaceae (3.70%) and Scrophulariaceae (<1%) families were determined at different percentages in the propolis sample. The pollen types found in the structure of propolis give information about the

flora of the region and also contribute to the understanding of the plants visited by bees. The results indicate that plant taxa belonging to Fabaceae, Lamiaceae and Asteraceae families are visited by bees for collecting nectar, propolis or pollen. These families are among the most common plant families in Turkey and their presence in propolis in large amounts is to be expected.

As a result of GC-MS analysis of propolis, benzyl alcohol, phenylethyl alcohol, benzoic acid, catechol, 4-vinyl-phenol, 2-propen-1-ol, 2-methoxy-4-vinylphenol, 4-hydroxy-3-methoxy, trans-cinnamic acid, gamma-murolene, delta-cadinene, naphthalene, tau-cadinol, alpha-copaene, t-murolol, (2e)-3-phenylpent-2,4-dienoic acid, benzyl benzoate, (2s,4as,5s,8ar)-perhydro-5,8a-dimethylnaphthalene-2-ol, 3,4-dimethoxycinnamic acid, benzyl cinnamate, p-coumaric acid, n1,n3-dimethyl-8-cyclohexylxanthin, eicosane, 2-propen-1-one, cinnamyl cinnamate, trifluoroacetic acid, 4H-1-benzopyran-4-one, 2,3-dihydro-5,7-d,hydroxy-2-phenyl-,(s), eicosane, 2-methoxymethyl-5-(4-methoxyphenyl)pyrrolo(1,2-c)pyrimidin-1(2h)-one, 4H-1-benzopyran-4-one, 5-hydroxy-7-methoxy-2-phenyl, 9,10-anthracenedione, 1,8-dihydroxy-3-methoxy-6-methyl-, diethyl 2-acetoxy-2-(2-oxocyclohexyl)malonate, chrysin, 5,7-dihydroxy-3-methoxy-2-phenyl-4H-1-benzopyran-4-one, estra-4,9,11-trien-3-one, 4',5-dihydroxy-7-methoxyflavanone, 9-nonadecene, docosyl pentafluoropropionate individual compounds were detected in different concentrations (Table 1). These compounds possess many biological properties. For instance, benzoic acid (RT/9.24) has antioxidant (24) and antimicrobial (25)


properties. Similarly, chrysin (RT/36.345), is a flavonoid that exhibits many biological activities and pharmacological effects, including anti-inflammatory, anticancer, antioxidant, and antiviral activities (26). This compound revealed a high peak area in our propolis, which may mean that propolis also exhibits the above-listed activities. In different studies, benzoic acid, 4-vinyl-phenol, eicosane, chrysin, delta-cadinene, phenylethyl alcohol, benzyl alcohol compounds were determined for propolis types from different origins (2, 27-29). The similarities observed in the samples of propolis from different origins suggest that the same plants are the source of these propolis samples.

Table 1. Chemical Screening of Propolis Composition

Retention Time	Compounds	% of total ion current
7.20	Benzyl alcohol	0.56
8.45	Phenylethyl alcohol	0.68
9.24	Benzoic acid	1.71
9.61	Catechol	0.15
9.91	4 - vinyl - phenol	0.54
11.25	2-Propen-1-ol	0.44
11.38	2-Methoxy-4-vinylphenol	1.86
12.80	4-hydroxy-3-methoxy	0.10
13.24	Trans-cinnamic acid	0.36
15.19	Gamma.-muurolene	0.19
15.34	Delta.-cadinene	0.23
15.66	Naphthalene	0.09
17.83	Tau.-cadinol	0.45
17.93	Alpha.-copaene	0.15
18.12	T-muurolol	0.92
19.12	(2E)-3-Phenylpent-2,4-dienoic acid	0.27
20.36	Benzyl benzoate	0.12
21.42	(2S,4as,5S,8ar)-perhydro-5,8a-dimethylnaphthalen-2-ol	0.42
23.21	3,4-Dimethoxycinnamic acid	0.51
26.71	Benzyl cinnamate	0.22
27.01	P-coumaric acid	0.13
28.77	N1,N3-Dimethyl-8-cyclohexylxanthin	2.38
30.22	Eicosane	0.43
31.99	2-Propen-1-one	7.61
32.12	Cinnamyl cinnamate	1.06
33.05	Trifluoroacetic acid	0.08
33.31	4H-1-benzopyran-4-one, 2,3-dihydro-5,7-d,hydroxy-2-phenyl-,(s)	15.35
33.42	Eicosane	0.31
34.76	2-Methoxymethyl-5-(4-methoxyphenyl)pyrrolo(1,2-c)pyrimidin-1(2H)-one	4.27
34.98	4H-1-Benzopyran-4-one, 5-hydroxy-7-methoxy-2-phenyl	5.26
35.53	9,10-Anthracenedione, 1,8-dihydroxy-3-methoxy-6-methyl-	0.31
36.18	Diethyl 2-acetoxy-2-(2-oxocyclohexyl)malonate	6.73
36.34	Chrysin	11.85
36.88	5,7-Dihydroxy-3-methoxy-2-phenyl-4H-1-benzopyran-4-one	1.18
37.00	Estra-4,9,11-trien-3-one	2.64
37.47	4',5'-Dihydroxy-7-methoxyflavanone	1.40
41.96	9-nonadecene	0.79
46.39	Docosyl pentafluoropropionate	0.75

In addition to the determination of the general chemical profile by GC-MS of the propolis extract, quantitative detection of the active ingredients of naringenin, pinocembrin and galangin was made using UHPLC (Figure 1). As a result of the investigation, the ratio of the active ingredients of the naringenin, pinocembrin and galangin of the extract was 2.45, 9.92, 7.06 mg/ml, respectively. Naringenin, pinocembrin and galangin are the essential phenolic compounds found in propolis extracts from Turkey and, hence, are most likely to contribute to the antioxidant activity of propolis extracts. These compounds were also reported by Guzelmeric et al. (2018) and Morlock et al. (2014) (30, 31). Also, Ristivojevi et al. (2014) identified pinocembrin, galangin, chrysin and CAPE as specific markers of *Populus nigra* buds (32). This could indicate that the resin source for the propolis used in the context of our work may be *Populus nigra*. Similarly, Yang & You (2017) reported that pinocembrin with its antioxidant, antimicrobial, vasorelaxant, anti-inflammatory and neuroprotective effects, etc. was present at the highest concentration in propolis (33). Naringenin, a flavonoid in the class of flavanones, can be extracted from various natural products. Different reports have shown the biological effects of naringenin, including blood lipid- and cholesterol-lowering effects (34), anti-inflammatory (35) and analgesic (35) activities (36). However, it is believed that the degree of such protective activities in propolis varies depending on the synergistic effects of individual compounds in its chemical composition.

Phenolic substances are important food components in terms of human health because of their effects on taste and odor formation, their participation

**Figure 1.** HPLC chromatogram of propolis extract

in color formation and change, their antimicrobial and antioxidative effects and their enzyme-inhibiting properties (37). Different types of bee products are assumed to be an effective source of natural antioxidants capable of resisting the impacts of oxidative stress underlying the pathogenesis of many diseases. Generally, compounds exhibiting phenolic characteristics, which express the ability to scavenge free radicals, are principally responsible for the antioxidant activity of bee products (38). For this reason, it is essential to determine the phenolic compositions of the products used as food or food supplements. There are several different techniques for the evaluation of the total phenolic, flavonoid or antioxidant capacity of synthetic antioxidants or bee products. However, the differences in these procedures make comparisons between studies in the literature difficult, and in some cases, conflicting results can be obtained. In this study, we used the Folin-Ciocalteu method and the aluminium chloride method to determine the total phenolic and flavonoid contents, respectively. These methods are assays that are widely used for phenolic profile determination in many different samples, including propolis (39, 40). In the literature, it has been reported that the antioxidant activity of multicomponent mixtures cannot be evaluated satisfactorily using a single antioxidant test due to many variables that may affect the results. Therefore, the application of several test procedures at the same time is recommended for better evaluation of antioxidant activity (41, 42). Herewith, the antioxidant activity of propolis was evaluated using different test methods together (CUPRAC, CERAC, ABTS) in our study. The CUPRAC method developed by Apak Güclu et al. (2004) was used to evaluate and confirm the total antioxidant capacity of biological samples (43). This method is based on the measurement of absorbance by the formation of a stable complex between neocuproine and copper (II) in the presence of neocuproine at 460 nm. The spectrophotometric CERAC assay is a method based on the determination of Ce (III) ions as a result of the reaction between Ce (IV)

ions and antioxidants in the sulfate acid medium. In the ABTS method, the preformed radical monocation of ABTS is produced by oxidation of ABTS with potassium persulfate and decreases in the presence of hydrogen-giving antioxidants. In this research, total phenolic and total flavonoid contents in propolis extract were determined 81.2±4 mg GAE/g and 164.8±8 mg QE/g of ethanol extract of propolis, respectively (Table 2). And all, antioxidant activity values of propolis were determined with a trolox equivalent of 282.8±9, 425.7±18, 186.4±8 mg TE/g according to the methods, CUPRAC, CERAC, and ABTS, respectively (Table 2). As in this study, Moreira, et al. (2008) reported that Portugal propolis is an important source of total phenol, with antioxidant properties that may be beneficial for human health (44). Ahn et al. (2004) obtained total phenolic and flavonoid content values ranging from 85-283 and 16-136 mg/g for ethanol extracts of propolis samples collected from different geographical regions (45). Kumazawa et al. (2004) determined the total phenolic and flavonoid contents of the propolis samples collected from different areas (Argentina, Brazil, China, New Zealand, South Africa, Thailand, Ukraine, United States, Uzbekistan) as 31.2-299 and 2.5-176 mg/g, respectively (46). The antioxidant activity of Algerian propolis was found to be greatly affected by TPC and TFC. This observation is consistent with previous studies of propolis. Differences in total phenolic and total flavonoid values are due to the fact that the phenolic content depends on geography, climate and plant origin (47).

The fatty acids composition of propolis extract is shown in Table 3. Saturated fatty acids were determined as capric acid, palmitic acid, stearic acid, arachidic acid, behenic acid and lignoceric acid. Monounsaturated fatty acids were detected as palmitoleic acid, oleic acid, cis-11-eicosenoic acid, erucic acid and nervonic acid. In addition, four polyunsaturated fatty acids were detected; linoleic acid, linolenic (ALA) acid, arachidonic acid, cis-13,16-docosadienoic acid. Similarly, it was reported that all propolis samples from Al-

Table 2. The results of bioactivity tests and extraction yield of propolis as mean±SD

	Extraction Yield (mg /g)	TPc (mg GAE /g)	TFc (mg QUE /g)	CUPRAC (mg TE /g)	CERAC (mg TE /g)	ABTS (mg TE /g)
Propolis extract	528.8±12	81.2±3	164.8±5	282.8±9	425.7±18	186.4±8

geria contained oleic, linoleic, stearic, eicosenoic, palmitoleic and palmitic acid, but only two samples did not contain arachidonic acid (3). The propolis oil was found to have cis-11-eicosenoic acid and erucic acid as its major fatty acid. cis-11-eicosenoic acid and erucic acid were present in amounts up to 31.87% and 22.93% respectively. cis-11-eicosenoic acid is beneficial as a raw material for medical supplies and a moisturizing component of cosmetic creams. It is also a precursor of erucic acid, which is beneficial for various applications such as cosmetics, creams, bio-diesel, lubricating oil, and therapeutic medicine (48). Studies on nervonic acid, another fatty acid that is highly (15.11%) detected in our propolis, showed a negative correlation between nervonic acid and obesity-related risk factors (49). This information suggests that our propolis sample has the potential to be used in these areas.

Conclusion

In conclusion, our study provides evidence that extract of propolis sampled from the western part of Istanbul, Turkey has high total phenolic content, and powerful antioxidant activity. Hence, it can be used as

a food supplement and a preventive agent for many diseases which are caused by free-radicals. Moreover, due to the specificity of total fatty acid composition, it can be used as a nutritional supplement as well as a potential raw material supply for the cosmetic sector.

Acknowledgments

This study was supported by Scientific Research Projects Unit (Project code: 26966) from Istanbul University.

References

- Huang S, Zhang C, Wang K, Li G, Hu F. Recent advances in the chemical composition of propolis. *Molecules* 2014; 19(12): 19610-32.
- Bayram N, Sorkun K, Oz G, Salih B, Topcu G. Chemical characterization of 64 propolis samples from Hakkari, Turkey. *Rec. Nat. Prod.* 2018; 12(6): 569-81.
- Rebiai A, Belfar M, Mesbahi M, Nani S, Tliba A, Amara DG, Chouikh A. Fatty acid composition of Algerian propolis. *J Fundam Appl Sci.* 2017; 9(3): 1656-71.
- Falcao S, Tomas A, Freire C, Vilas-Boas M. A voltammetric tool for the evaluation of propolis antioxidant activity. *Eur Food Res Technol.* 2016, 242: 1393-401
- Neto M, Tintino S, da Silva A, Costa M, Boligon A, Matias E, Balbino V, Menezes I, Coutinho H. Seasonal variation of Brazilian red propolis: Antibacterial activity, synergistic effect and phytochemical screening. *Food Chem Toxicol.* 2017; 107: 572-80.
- Freires I, Queiroz V, Furletti V, Ikegaki M, de Alencar S, Duarte M, Rosalen P. Chemical composition and antifungal potential of Brazilian propolis against *Candida* spp. *J Mycol Med.* 2016; 26(2): 122-32.
- Kujungiev A, Tsvetkova I, Serkedjieva Y, Bankova V, Christov R, Popov S. Antibacterial, antifungal and antiviral activity of propolis of different geographic origin. *J Ethnopharmacol.* 1999; 64(3): 235-40.
- Dobrowolski J, Vohora S, Sharma K, Shah S, Naqvi S, Dandiya P. Antibacterial, antifungal, antiamebic, anti-inflammatory and antipyretic studies on propolis bee products. *J Ethnopharmacol.* 1991; 35(1): 77-82.
- Ledon N, Casaco A, Gonzalez R, Merino N, Gonzalez A, Tolon Z. Antipsoriatic, anti-inflammatory, and analgesic effects of an extract of red propolis. *Acta Pharmacol. Sin.* 1997; 18(3): 274-6.
- De Campos Ro, Paulino N, Da Silva CH, Scremm A, Calixto JB. Natural Products: Anti-hyperalgesic Effect of an Ethanolic Extract of Propolis in Mice and Rats. *J. Pharm Pharmacol.* 1998; 50(10): 1187-93.

Table 3. Fatty acid composition of propolis sample (as a percentage of total fatty acids)

Retention Time	Common Name	Fatty acid	% of total
6.11	Capric acid	C10:0	0.33
10.37	Palmitic acid	C16:0	1.23
10.71	Palmitoleic acid	C16:1	0.4
12.77	Stearic acid	C18:0	0.39
13.10	Oleic acid	Cis- C18:1	1.29
13.77	Linoleic acid	Cis-C18:2	0.29
14.64	Linolenic (ALA) acid	C18:3n3	0.77
15.57	Arachidic acid	C20:0	1.23
15.94	cis-11-Eicosenoic acid	C20:1n9	31.87
17.38	Arachidonic acid	C20:4n6	0.93
18.5	Behenic acid	C22:0	1.19
18.9	Erucic acid	C22:1	22.93
19.70	Cis-13,16-Docosadienoic acid	C22:2	0.57
21.45	Lignoceric acid	C24:0	0.15
21.70	Nervonic acid	C24:1	15.14

11. Bayram NE, Karadayı M, Güllüce M, Bayram S, Sorkun K, Aydoğan MN, Koç TY, Alaylar B and Salih B. Genotoxic and antigenotoxic evaluation of propolis by using in vitro bacterial assay systems. *Mellifera*. 2015; 15(1): 29-36.
12. Mohammadzadeh S, Sharriatpanahi M, Hamed M, Amanzadeh Y, Ebrahimi S, Ostad S. Antioxidant power of Iranian propolis extract. *Food Chem*. 2007; 103(3): 729-33.
13. Ecem-Bayram N, Gercek Y. Appropriate maceration duration for the extraction of propolis. *Fresen Environ Bull*. 2019; 28(1): 188-92.
14. Gajger I, Pavlovic I, Bojic M, Kosalec I, Srecec S, Vlainic T, Vlainic J. The components responsible for the antimicrobial activity of propolis from continental and mediterranean regions in Croatia Czech J. Food Sci. 2017; 35(5): 376-85.
15. Warakomska Z, Maciejewicz W. Microscopic analysis of propolis from polish regions. *Apidologie* 1992; 23(4): 277-83.
16. Ecem Bayram N, Sorkun K, Cevahir Oz G. Plant origins of propolis from Hakkari, Turkey. *GU J Sci*. 2018; 31(3): 725-38.
17. Magalhaes L, Santos F, Segundo M, Reis S, Lima J. Rapid microplate high-throughput methodology for assessment of Folin-Ciocalteu reducing capacity. *Talanta*. 2010; 83: 441-7.
18. Jia Z, Tang M, Wu J. The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chem*. 1999; 64(4): 555-9.
19. Apak R, Guclu K, Demirata B, Ozyurek M, Celik S, Bektasoglu B, Berker K, Özyurt D.. Comparative evaluation of various total antioxidant capacity assays applied to phenolic compounds with the CUPRAC assay. *Molecules*. 2007; 12(7): 1496-547.
20. Ozyurt D, Demirata B, Apak R. Determination of total antioxidant capacity by a new spectrophotometric method based on Ce (IV) reducing capacity measurement. *Talanta*. 2007; 71(3): 1155-65.
21. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med*. 1999; 26(9-10): 1231-7.
22. Regulation H. Commission Regulation (EEC) No. 2568/91 of 11 July 1991 on the characteristics of olive oil and olive-residue oil and on the relevant methods of analysis Official Journal L 248, 5 September 1991. *Offic J L* 1991, 248: 1-83.
23. IUPAC. Standard methods for the analysis of oils, fats and derivatives. International union of pure and applied chemistry 1987.
24. Sz wajgier D, Pielecki J, Targoński Z. Antioxidant activities of cinnamic and benzoic acid derivatives. *Acta Sci Pol Technol Aliment..* 2005; 4(2): 129-42.
25. Park E, Moon W, Song M, Kim M, Chung K, Yoon J. Antimicrobial activity of phenol and benzoic acid derivatives. *Int Biodeterior Biodegradation*. 2001, 47: 209-14.
26. Mani R, Natesan V. Chrysin: Sources, beneficial pharmacological activities, and molecular mechanism of action. *Phytochemistry* 2018, 145: 187-96.
27. Bayram NE. Major Constituents of Different Propolis Samples. *Hacettepe J. Biol. & Chem*. 2018; 45(4): 581-4.
28. Bayram S, Bayram NE, Gercek YC, Aydoğan MN, Oz GC. Chemical analysis and antimicrobial effect of propolis from Hakkari Province of Turkey against some pathogenic microorganisms. *Eur J Biol* 2017; 76(2): 74-8.
29. Silici S, Kutluca S. Chemical composition and antibacterial activity of propolis collected by three different races of honeybees in the same region. *J Ethnopharmacol*. 2005; 99(1): 69-73.
30. Guzelmeric E, Ristivojevic P, Trifkovic J, Dastan T, Yilmaz O, Cengiz O, Yesilada E. Authentication of Turkish propolis through HPTLC fingerprints combined with multivariate analysis and palynological data and their comparative antioxidant activity. *LWT-Food Sci Technol* . 2018; 87: 23-32.
31. Morlock G, Ristivojevic P, Chernetsova E. Combined multivariate data analysis of high-performance thin-layer chromatography fingerprints and direct analysis in real time mass spectra for profiling of natural products like propolis. *J Chromatogr*. 2014; 1328: 104-12
32. Ristivojevic P, Andric F, Trifkovic J, Vovk I, Stanisavljevic L, Tesic Z, Milojkovic-Opsenica D. Pattern recognition methods and multivariate image analysis in HPTLC fingerprinting of propolis extracts. *J Chemom*. 2014; 28(4): 301-310.
33. Yang Z, You Y. P19 Intracerebral microdialysis studies on the role of pinocembrin on CYP3A in rat. *Biochem Pharmacol*. 2017; 139: 131.
34. Santos F, Bastos E, Uzeda M, Carvalho M, Farias L, Moreira E. Antibacterial activity of propolis produced in Brazil against *Actinobacillus actinomycetemcomitans*, *Fusobacterium* spp. and bacteria from the *Bacteroides fragilis* group isolated from human and marmoset hosts. *Anaerobe*. 1999; 5(3-4): 479.
35. Manchope M, Casagrande R, Verri W. Naringenin: an analgesic and anti-inflammatory citrus flavanone. *Oncotarget*. 2017; 8(3): 3766-7.
36. Sangpheak W, Kicuntod J, Schuster R, Rungrotmongkol T, Wolschann P, Kungwan N, Viernstein H, *et al*. Physical properties and biological activities of hesperetin and naringenin in complex with methylated β -cyclodextrin. *Beilstein J Org Chem*. 2015; 11(1): 2763-73.
37. Kolaç T, Gürbüz P, Yetiş G. Doğal ürünlerin fenolik içeriği ve antioksidan özellikleri. İnönü Üniversitesi Sağlık Hizmetleri Meslek Yüksek Okulu Dergisi 2017; 5(1): 26-42.
38. Kocot J, Kielczykowska M, Luchowska-Kocot D, Kurzepa J, Musik I. Antioxidant potential of propolis, bee pollen, and royal jelly: Possible medical application. *Oxid Med Cell Longev*. 2018; 1-29.
39. Popova M, Giannopoulou E, Skalicka-Wozniak K, Graikou K, Widelski J, Bankova V, Kalofonos H, Sivolapenko K, Gawel-Beben B, Antosiewicz, Chinou I. Characterization and biological evaluation of propolis from Poland. *Molecules*. 2017; 22(7):1-13.
40. Christina D, Hermansyah H, Wijanarko A, Rohmatin E, Sahlan M, Pratami D, Im A, *et al*. Selection of Propolis *Tetragonula* sp. extract solvent with flavonoids and polyphenols concentration and antioxidant activity parameters. 2nd

- Biomedical Engineerings Recent Progress in Biomaterials, Drugs Development, and Medical Devices 2018, 1933.
41. Antolovich M, Prenzler P, Patsalides E, McDonald S, Roberts K. Methods for testing antioxidant activity. *Analyst*. 2002; 127(1): 183-98.
 42. Capanoglu E, Beekwilder J, Boyacioglu D, Hall R, De Vos R. Changes in antioxidant and metabolite profiles during production of tomato paste. *Agric Food Chem*. 2008; 56(3): 964-73.
 43. Apak R, Guclu K, Ozyurek M, Karademir S. Novel total antioxidant capacity index for dietary polyphenols and vitamins C and E, using their cupric ion reducing capability in the presence of neocuproine: CUPRAC method. *J Agric Food Chem*. 2004, 52(26): 7970-81.
 44. Moreira L, Dias L, Pereira J, Estevinho L. Antioxidant properties, total phenols and pollen analysis of propolis samples from Portugal. *Food Chem Toxicol*. 2008, 46(1): 3482-85.
 45. Ahn M, Kumazawa S, Hamasaka T, Bang K, Nakayama T. Antioxidant activity and constituents of propolis collected in various areas of Korea. *J Agric Food Chem*. 2004; 52(24): 7286-92.
 46. Kumazawa S, Murase M, Momose N, Fukumoto S. Analysis of antioxidant prenylflavonoids in different parts of *Macaranga tanarius*, the plant origin of Okinawan propolis *Asian Pac J Trop Med*. 2014; 7(1): 16-20.
 47. Choi Y, Noh D, Cho S, Suh H, Kim K, Kim J. Antioxidant and antimicrobial activities of propolis from several regions of Korea. *LWT-Food Sci Technol*. 2006; 39(7): 756-61.
 48. Kikukawa H, Sakuradani E, Nishibaba Y, Okuda T, Ando A, Shima J, Shimizu S, Ogawa J. Production of cis-11-eicosenoic acid by *Mortierella* fungi. *J Appl Microbiol*. 2015; 118(3): 641-7.
 49. Oda E, Hatada K, Kimura J, Aizawa Y, Thanikachalam P, Watanabe K. Relationships between serum unsaturated fatty acids and coronary risk factors - Negative relations between nervonic acid and obesity-related risk factors *Int Heart J*. 2005; 46(6): 975-85.
-
- Correspondence:
Nesrin Ecem Bayram
Department of Food Processing, Aydintepe Vocational School,
Bayburt University, Bayburt 69000, Turkey,
Fax: +904583332028
E-mail address: ecem.nesrin@gmail.com